

Annual Report

North Dakota Air Quality Monitoring Data Summary 2001



Annual Report

North Dakota

Air Quality Monitoring Data Summary 2001

March 2002

John Hoeven
Governor

Terry L. Dwelle, MD
State Health Officer

Francis J. Schwindt
Environmental Health Section Chief



North Dakota Department of Health
Division of Air Quality
Air Quality Monitoring Branch
1200 Missouri Ave.
Bismarck, N.D. 58506-5520

TABLE OF CONTENTS

LIST OF TABLES	iii
LIST OF FIGURES	v
LIST OF APPENDICES	vii
EXECUTIVE SUMMARY	1
INTRODUCTION	3
DESCRIPTION	4
Department Sites	4
Industry Sites	4
NETWORK CHANGES	8
Department Changes	8
Industry Changes	8
MONITORING RESULTS	9
Introduction	9
Sulfur Dioxide	11
Sulfur Dioxide 5-Minute Average	14
Nitrogen Dioxide	16
Ammonia	18
Ozone	19
Particulate Matter (PM _{2.5} & PM ₁₀)	22
Inhalable PM _{2.5} Particulates	23
Inhalable Continuous PM _{2.5} Particulates	24
Inhalable PM ₁₀ Particulates	25
SUMMARY AND CONCLUSIONS	27
REFERENCES	29
APPENDICES	33

LIST OF TABLES

<u>Table No.</u>		<u>Page No.</u>
1	State AAQM Network Description	6
2	Sulfur Dioxide	13
3	SO ₂ 5-Minute Averages	15
4	Nitrogen Dioxide	17
5	Ammonia	18
6	Ozone	21
7	Inhalable PM _{2.5} Particulates	24
8	Inhalable Continuous PM _{2.5}	25
9	Inhalable PM ₁₀ Particulates	25
A1-1	North Dakota Ambient Air Quality Standards	37
A1-2	Federal Ambient Air Quality Standards	38

LIST OF FIGURES

<u>Figure No.</u>		<u>Page No.</u>
1	North Dakota Air Quality Monitoring Network	7
A2-1	Environmental Engineering Organizational Chart	41
A3-1	Amerada Hess Star Charts	45
A3-2	Beulah Star Charts	46
A3-3	Bear Paw Star Charts	47
A3-4	Dunn Center Star Charts	48
A3-5	DGC Star Charts	49
A3-6	Fargo Star Charts	51
A3-7	Hannover Star Charts	52
A3-8	Mandan/Mandan NW Star Charts	53
A3-9	TRNP - NU Star Charts	54
A3-10	Short Creek - SPM/TRNP - NU	55
A4-1	Amerada Hess/Beulah North Trends	59
A4-2	Beulah North (cont.)/Bear Paw Trends	60
A4-3	DGC Trends	61
A4-4	Dunn Center/Dickinson/Dragswolf Trends	62
A4-5	Fargo NW Trends	63
A4-6	Grand Forks - North/Hannover Trends	64
A4-7	Mandan/Mandan NW Trends	65
A4-8	TRNP - NU Trends	66
A4-9	TRNP - SU Trends	67

LIST OF APPENDICES

<u>Appendix No.</u>		<u>Page No.</u>
1	North Dakota and Federal Ambient Air Quality Standards	35
2	Air Quality Personnel Organizational Chart	39
3	Wind and Pollution Star Charts	43
4	1992-2001 Trends	57

EXECUTIVE SUMMARY

The North Dakota Department of Health operated ten ambient and two special purpose air quality monitoring sites, industry operated eight source-specific air quality monitoring sites and the Three Affiliated Tribes on the Fort Berthold Indian Reservation operated two ambient sites. The data from these sites indicated that the quality of the ambient air in North Dakota was generally good during 2001.

There were no sulfur dioxide, nitrogen dioxide, ozone or particulate matter exceedances of either the state or federal ambient air quality standards measured during the year. Through legislative action effective August 1, 1997, coal conversion facilities and oil refineries were exempted from the state sulfur dioxide standards. Therefore, any values listed as an exceedance in the data summaries are subject to further manual review to determine the most likely source(s) causing the listed exceedance.

INTRODUCTION

The North Dakota Department of Health, Environmental Health Section, Division of Air Quality, henceforth known as the department, has the primary responsibility for protecting the health and welfare of North Dakotans from the harmful effects of air pollution. The department ensures that the ambient air quality in North Dakota is better than the levels required by the state and federal Ambient Air Quality Standards^{1,2} and the "Prevention of Significant Deterioration of Air Quality Rules."³ To address this responsibility, the department operates a network of ambient air quality monitors.

In addition to the state-operated ambient air quality monitoring sites, three industrial sources of air pollutants operated air quality monitoring sites within their immediate spheres of influence. These site locations are selected based on computer dispersion modeling and prevailing wind directions.

This report provides an overview of air quality monitoring activities conducted by the department and industry during the 12-month period beginning Jan. 1, 2001, and ending Dec. 31, 2001. The report includes data summaries for the monitored pollutants and significant changes that occurred to the monitoring program. Also included are wind and pollution star charts and trend graphs. The pollution star charts (Appendix 3) indicate the percentage of time a pollutant is detected when the wind is from each direction. The trend graphs (Appendix 4) show the maximum concentration for each pollutant standard and the percentage of time a concentration was above the minimum detectable limit for the specific analysis method.

The Three Affiliated Tribes on the Fort Berthold Indian Reservation operate a tribal network that consists of two sites: White Shield and Drags Wolf. The data summaries are included only for informational purposes, since tribal data is not subject to state ambient air quality standards.

NETWORK DESCRIPTION

Department Sites

During 2001, the department operated twelve air quality monitoring sites. Ten were ambient monitoring sites, and two were special purpose monitoring (SPM) sites near the AMOCO Refinery and MDU Heskett Power Plant at Mandan. Table 1 lists the department monitoring sites which were active during the year.

In general, department ambient air quality monitoring (AAQM) sites obtain air quality data to meet five objectives: (1) determine representative concentrations in areas of high population density (urban or population oriented monitoring), (2) determine general background concentration levels, (3) measure highest concentrations expected to occur in an area covered by an individual site, (4) determine representative impacts on ambient air quality levels near significant sources, and , (5) determine the effects of long-range pollution transport.

The department's ambient air quality monitoring network normally does not include source-specific monitoring; i.e., monitoring a single, specific source. However, the two Mandan SPM sites were established to collect source-specific 5-minute peak and hourly sulfur dioxide averages due to impacts from the MDU Heskett Power Plant and AMOCO Refinery.

The department is working with Environment Canada, the Environmental Protection Agency (EPA), Saskatchewan Environment and Resource Management (SERM) and SASKPower to operate a North Dakota-Saskatchewan Transboundary ambient air quality monitoring network with three sites (Rafferty Dam, Estevan, Boundary Dam Power Station) in Saskatchewan and the two sites (Short Creek and Lignite) in North Dakota. The ND-SK Transboundary network became fully operational Dec. 5, 2000, when the Estevan PM_{2.5} sampler collected its first sample. Data collected at these five sites are addressed in that network's own annual reports.

The department, in issuing Permits to Construct and Permits to Operate for major sources, may require those sources to operate ambient air quality monitoring programs to assess impacts on local air quality.

Industry Sites

Industry operated eight source-specific air quality monitoring sites during the year. Table 1 also lists the industry networks and monitoring sites active during the year.

In general, industry air quality monitoring sites obtain data at locations expected to show high concentrations of pollution from a specific source. These source-specific sites are selected using computer dispersion modeling programs and annual wind patterns. The distance a monitoring site is located from a source is determined by the primary pollutant monitored.

Figure 1 displays both department and industry monitoring sites. If an industry has more than one site, only the general location within the county is indicated. This principle also applies to the Mandan location, which represents the two sites at Mandan.

TABLE 1

State AAQM Network Description

Site Name AQS Site #	Type Station	Parameter Monitored ¹	Operating Schedule	Monitoring Objective ²	Spatial Scale ²	Date Site Began
1 Beulah North	SLAMS	PM _{2.5} SO ₂ , NO ₂ , O ₃ , MET NH ₃ cont. PM _{2.5}	6 th Day cont. cont. cont.	Population Exposure Population Exposure Population Exposure Population Exposure	Neighborhood Neighborhood Regional Neighborhood	12/98 04/80 11/00 10/00
2 Bismarck Residential	SLAMS	PM ₁₀ PM _{2.5} PM _{2.5} Speciation	6 th Day 3 rd Day 3 rd Day	Population Exposure Population Exposure Long-range Transport	Urban Urban Regional	1/01 12/98 1/01
3 Dickinson Residential ³	SLAMS	PM _{2.5}	6 th Day	Population Exposure	Urban	07/89
4 Drags Wolf	Tribal	MET PM ₁₀	cont. 6 th Day	General Background	Regional	01/86
5 Dunn Center	SLAMS	SO ₂ , NO ₂ , O ₃ , MET	cont.	General Background	Regional	10/79
6 Fargo NW	SLAMS	PM _{2.5} PM _{2.5} SO ₂ , NO ₂ , O ₃ , MET cont. PM _{2.5} PM _{2.5} Speciation	3 rd day 3 rd Day cont. cont. 3 rd Day	Population Exposure Collocated Population Exposure Population Exposure Long-range Transport	Urban N/A Urban Urban Regional	12/98 05/98 7/00 7/01
7 Grand Forks North ⁴	SLAMS	PM _{2.5}	3 rd Day	Population Exposure	Urban	12/98
8 Hannover	SLAMS	SO ₂ , NO ₂ , O ₃ , MET	cont.	General Background	Regional	10/84
9 Mandan Refinery - SPM	SPM	SO ₂ , MET	cont.	Source Impact	Neighborhood	12/95
10 Mandan Refinery NW - SPM	SPM	SO ₂ , MET	cont.	Source Impact	Neighborhood	09/98
11 Sharon ⁴	SLAMS	PM _{2.5}	6 th Day	General Background	Regional	12/98
12 TRNP - NU	SLAMS	SO ₂ , NO ₂ , O ₃ , MET	Cont.	Source Impact	Regional	02/99
13 TRNP - SU	SLAMS	SO ₂ , O ₃ MET PM _{2.5}	cont. 6 th Day	General Background	Regional	07/95 7/00
14 White Shield	Tribal	SO ₂ , MET PM ₁₀	cont. 6 th Day	General Background	Regional	07/90
Company	Site Name					
15 Amerada Hess Corporation	TIOGA #1 TIOGA #3	SO ₂ , MET SO ₂	cont. cont.	Source Impact Source Impact	Urban Urban	07/87 11/87
16 Bear Paw Energy, Inc.	MGP #3 MGP #5	SO ₂ , MET SO ₂ , MET	cont. cont.	Source Impact Source Impact	Urban Urban	11/94 01/98
17 Dakota Gasification Company	DGC #12 DGC #14 DGC #16 DGC #17	SO ₂ , NO ₂ , MET SO ₂ SO ₂ SO ₂ , NO ₂	cont. cont. cont. cont.	Source Impact Source Impact Source Impact Source Impact	Urban Urban Urban Urban	01/80 01/89 10/95 10/95
1. MET refers to meteorological and indicates wind speed and wind direction monitoring equipment. 2. Not applicable to MET. 3. Terminated June 30. 4. Terminated December 31.						




Figure 1. North Dakota Air Quality Monitoring Network

NETWORK CHANGES

Department Changes

Changes to the state monitoring network consisted of adding, terminating and moving the following samplers to new locations. Speciation samplers were added to Bismarck Residential and Fargo NW along with particulate matter (<10 microns) (PM_{10}) samplers. The Theodore Roosevelt National Park - North Unit (TRNP - NU) site was re-activated in August 2001 with SO_2 , NO_2 , O_3 , and MET. The Grand Forks and Sharon $PM_{2.5}$ samplers were terminated effective Dec. 31, 2001 with both of these samplers moved to TRNP - NU.

The Fargo NW speciation sampler was installed at EPA's request and is a part of the EPA National Trends Network. As a part of preparing for the regional haze/long-range transport rule, the sulfate and nitrate concentrations were modeled using the latest dispersion modeling techniques. The result of the modeling indicated the Bismarck Residential site was located midway between the maximum sulfate (SO_4^{2-}) and nitrate (NO_3^-) areas. Therefore, the Bismarck Residential site was scheduled to receive a speciation sampler. To address the need for long-range transport/regional haze monitoring within the state network, TRNP - NU was re-activated and is scheduled to receive a speciation sampler effective Jan. 1, 2002.

Along with each speciation sampler, a Federal Reference Method $PM_{2.5}$ and Federal Equivalent Method PM_{10} sampler will be operated. The rationale for this monitoring arrangement is to collect the same data set as collected by the IMPROVE (Interagency Monitoring of PROtected Visual Environments) samplers installed at Theodore Roosevelt National Park - South Unit (TRNP - SU) and Lostwood National Wildlife Refuge (NWR).

Industry Changes

No changes were made to the industry networks.

MONITORING RESULTS

Introduction

Ambient and source-specific air quality data collected during the year at monitoring sites operated by the department and industry are summarized in tables for the following pollutants: sulfur dioxide (SO_2), nitrogen dioxide (NO_2), ozone (O_3), ammonia (NH_3), federal reference method inhalable fine particulates ($PM_{2.5}$), and inhalable coarse particulates (PM_{10}). Each section contains a description of the physical characteristics and health effects, a comparison to the state standards, and a data summary.

The data summaries for gaseous pollutants include maximum concentrations, month/day/hour of each maximum, arithmetic means and the percentage of readings greater than the minimum detectable value (MDV) for the analytical method used for each parameter. Where applicable, the number of times a state standard was exceeded is indicated. The concentrations for gaseous pollutants are reported in parts per billion (ppb).

The $PM_{2.5}$ and PM_{10} data summaries contain the three highest 24-hour average concentrations; month/day of each maxima; annual arithmetic mean; the number of times the 24-hour standard was exceeded, if applicable; and an asterisk (*) if the annual standard is exceeded, if applicable. The concentrations are reported in micrograms per cubic meter ($\mu\text{g}/\text{m}^3$).

Continuous $PM_{2.5}$ data summaries contain the two highest 1-hour averages and the four highest 24-hour averages; the annual average; the number of times the 24-hour standard was exceeded; if applicable, and an asterisk (*) if the annual standard is exceeded, if applicable. The concentrations are reported in micrograms per cubic meter ($\mu\text{g}/\text{m}^3$).

For statistical purposes, pollutant concentrations less than the minimum detectable value (MDV) for the analytical method used are assigned a value equal to one-half the MDV. The MDV for SO_2 and NO_2 is 2 ppb; O_3 is 4 ppb; $PM_{2.5}$ is $2.0 \mu\text{g}/\text{m}^3$; and PM_{10} is $4 \mu\text{g}/\text{m}^3$. The MDV for the continuous $PM_{2.5}$ is $-2.0 \mu\text{g}/\text{m}^3$. Annual means are calculated for SO_2 , NO_2 , $PM_{2.5}$, and PM_{10} . However, only those means with more than 75 percent of data greater than the MDV are unbiased calculations.

As part of the statistical evaluation, the data recovery (NUM OBS) is evaluated to determine if the data recovery complies with the state's required 80 percent data recovery rate. A continuous analyzer operating fewer than 7,028 hours per year may achieve at least an 80 percent data recovery for the period operated; However, it does not meet the 80 percent data recovery for the full year. Each analyzer at a site not meeting the 80 percent data recovery for the year is flagged in the "NUM OBS" column by

placing “***” underneath the number of observations. Particulate matter samplers must collect at least 48 samples per year for 1-in-6 day sampling and 96 samples per year for 1-in-3 day sampling to meet the 80 percent data recovery rate.

Sulfur Dioxide

Physical Characteristics and Sources

Sulfur dioxide is a colorless gas with a pungent odor detectable by the human nose at concentrations of 500 to 800 ppb.⁴ It is highly soluble in water where it forms sulfurous acid (H_2SO_3). In the atmosphere, sulfurous acid is easily converted to sulfuric acid (H_2SO_4), the major acidic component of “acid rain,” which then may convert to a sulfate. On a worldwide basis, sulfur dioxide is considered to be a major pollutant. It is emitted mainly from stationary sources that burn coal and oil – such as utility boilers. Other sources of sulfur dioxide include refineries, natural gas processing plants, oil well heaters and flares.

Health Effects

Sulfur dioxide can be converted in the atmosphere to sulfuric acid aerosols and particulate sulfate compounds which are corrosive and potentially carcinogenic (cancer-causing). The major health effects of sulfur dioxide appear when it is associated with high levels of other pollutants, such as particulate. Sulfur dioxide also may play an important role in the aggravation of chronic illnesses, such as asthma. The incidence and intensity of asthma attacks have increased when asthmatics are exposed to higher levels of sulfur dioxide and particulate matter sulfates, which are products of atmospheric sulfur dioxide reactions.⁴

Standards Comparison

Sulfur dioxide was monitored at 17 sites. Eight sites were run by the department, eight by industry, and one by the Three Affiliated Tribes on the Fort Berthold Indian Reservation. As a result of legislative action effective Aug. 1, 1997, coal conversion facilities and oil refineries were exempted from the state sulfur dioxide standards leaving these two classes of sources subject only to the federal standards. Therefore, the Dakota Gasification Company (DGC) network, Mandan NW - SPM, and Mandan - SPM are compared only to the federal standards.

The 1-hour state standard (273 ppb) was not exceeded during the year by an applicable source. The maximum 1-hour concentration was 164 ppb at DGC #12.

The 3-hour federal secondary standard (500 ppb) was not exceeded during the year. The maximum 3-hour average concentration was 116 ppb at Mandan - SPM.

The 24-hour state standard (99 ppb) was not exceeded twice during the year. The maximum 24-hour average concentration was 38 ppb at Mandan - SPM. .

Among those sites that collected at least 80 percent of the possible data during the year, the maximum annual arithmetic mean was 5.5 ppb at Mandan - SPM.

The sulfur dioxide data are summarized in Table 2.

TABLE 2

COMPARISON OF AIR QUALITY DATA WITH
THE NORTH DAKOTA AMBIENT AIR QUALITY STANDARDS *

POLLUTANT : Sulfur Dioxide (ppb)

LOCATION	YEAR	SAMPLING PERIOD	NUM OBS	1 - HOUR		M A X 3 - HOUR		24 - HOUR		ARITH MEAN	1HR #>273	24HR #>99	% >MDV
				1ST MM/DD/HH	2ND MM/DD/HH	1ST MM/DD/HH	2ND MM/DD/HH	1ST MM/DD	2ND MM/DD				
Amerada Hess - Tioga #1	2001	JAN-DEC	8192	57 06/10/18	57 11/19/22	30 11/19/23	29 11/08/11	7 11/19	6 11/08	1.4			12.1
Amerada Hess - Tioga #3	2001	JAN-DEC	8634	81 11/18/09	76 06/18/17	64 06/18/17	50 11/18/11	22 05/21	20 03/24	2.9			24.8
Bear Paw - MGP #3	2001	JAN-DEC	8695	48 08/30/07	21 02/05/10	18 08/30/08	11 12/12/08	3 12/12	3 08/30	1.2			8.0
Bear Paw - MGP #5	2001	JAN-DEC	8690	47 02/10/08	28 08/18/07	18 02/10/08	13 02/10/11	6 02/10	3 06/03	1.3			10.8
Beulah - North	2001	JAN-DEC	8691	89 07/06/07	87 03/28/13	67 03/28/14	66 03/24/11	26 03/24	17 03/28	2.0			22.7
DGC #12	2001	JAN-DEC	8686	164 03/12/22	108 03/12/19	99 03/12/23	96 03/12/20	27 03/12	14 03/28	2.2			31.4
DGC #14	2001	JAN-DEC	8707	111 03/15/10	95 01/17/10	65 03/15/14	63 03/15/11	17 03/15	11 03/13	1.9			20.7
DGC #16	2001	JAN-DEC	8570	96 08/08/07	75 08/07/12	47 08/07/14	42 03/16/14	10 08/07	10 08/18	2.1			21.6
DGC #17	2001	JAN-DEC	8687	84 08/07/12	72 08/07/13	54 08/07/14	48 08/08/11	13 08/07	11 08/08	1.9			19.6
Dunn Center	2001	JAN-DEC	8360	25 09/15/09	20 07/09/08	14 07/24/11	13 09/15/11	5 02/14	5 07/09	1.3			12.5
Fargo NW	2001	JAN-DEC	8372	12 02/11/05	9 02/27/23	7 02/27/20	7 02/27/23	4 02/27	3 02/11	1.1			4.0
Hannover	2001	JAN-DEC	5465 ***	85 01/24/03	62 03/25/09	51 03/25/11	50 01/24/05	16 01/24	10 03/25	2.1			21.1
Mandan - SPM	2001	JAN-DEC	8695	140 03/28/12	119 03/28/13	116 03/28/14	77 11/07/20	38 01/15	33 02/13	5.5			37.7
Mandan NW - SPM	2001	JAN-DEC	8661	115 09/27/10	91 09/27/09	90 09/27/11	74 12/14/20	26 09/25	23 03/28	3.9			42.4
TRNP - NU	2001	AUG-DEC	3637 ***	9 10/01/10	7 09/15/16	6 10/01/11	6 12/11/14	2 12/11	2 10/01	1.1			3.6
TRNP - SU (Painted Canyon)	2001	JAN-DEC	8700	20 02/07/17	16 07/17/19	15 02/07/17	12 07/17/20	4 02/07	3 07/17	1.1			6.7
White Shield	2001	JAN-DEC	8458	40 03/26/10	40 08/03/13	34 03/22/20	30 03/26/14	11 03/26	8 03/22	1.5			14.4

The maximum 1-hour concentration is 164 ppb at DGC #12 on 03/12/22

The maximum 3-hour concentration is 116 ppb at Mandan - SPM on 03/28/14

The maximum 24-hour concentration is 38 ppb at Mandan - SPM on 01/15

* The air quality standards are:

STATE Standards -

- 1) 273 ppb maximum 1-hour average concentration.
- 2) 99 ppb maximum 24-hour average concentration.
- 3) 23 ppb maximum annual arithmetic mean concentration.

FEDERAL Standards -

- 1) 500 ppb maximum 3-hour concentration not to be exceeded more than once per year.
- 2) 140 ppb maximum 24-hour concentration not to be exceeded more than once per year.
- 3) 30 ppb annual arithmetic mean.

*** Less than 80% of the possible samples (data) were collected.

Sulfur Dioxide 5-Minute Average

Sulfur dioxide 5-minute averages were collected at state-operated sites and the Bear Paw Energy network. The maximum 5-minute average was 229 ppb at Mandan - SPM.

The sulfur dioxide 5-minute data is presented in Table 3.

TABLE 3

COMPARISON OF AIR QUALITY DATA WITH
THE NORTH DAKOTA AMBIENT AIR QUALITY STANDARDS *POLLUTANT : SO₂ 5-Minute Averages (ppb)

POLLUTANT : SO ₂ 5-Minute Averages (ppb)												
LOCATION	YEAR	SAMPLING PERIOD	NUM OBS	1ST	5 - M I N U T E		M A X I M A		# HOURS >600	% >MDV		
					DATE MM/DD/HH	2ND DATE MM/DD/HH	3RD DATE MM/DD/HH					
Bear Paw - MGP #3	2001	JAN-DEC	8695	143	08/30/07	58	02/05/10	38	03/20/08	0	16.6	
Bear Paw - MGP #5	2001	JAN-DEC	8690	91	05/14/22	79	02/10/08	58	02/10/09	0	23.2	
Beulah - North	2001	JAN-DEC	8691	169	03/28/14	161	07/06/07	152	03/28/13	0	36.5	
Dunn Center	2001	JAN-DEC	8358	30	09/15/09	28	09/03/08	27	07/24/08	0	19.8	
Fargo NW	2001	JAN-DEC	8372	12	02/11/05	11	12/31/14	11	12/31/15	0	4.8	
Hannover	2001	JAN-DEC	5465 ***	173	01/24/03	118	01/24/02	93	03/25/10	0	30.8	
Mandan - SPM	2001	JAN-DEC	8695	229	03/28/13	225	02/03/20	203	09/30/11	0	49.3	
Mandan NW - SPM	2001	JAN-DEC	8661	222	01/08/14	173	09/27/10	153	09/27/09	0	56.8	
TRNP - NU	2001	AUG-DEC	3637 ***	12	10/01/09	12	10/01/10	10	10/01/11	0	5.7	
TRNP - SU (Painted Canyon)	2001	JAN-DEC	8700	20	02/07/17	17	11/11/23	16	07/17/19	0	8.6	

The maximum 5-minute concentration is 229 ppb at Mandan - SPM on 03/28/13

* No Standard is currently in effect.

*** Less than 80% of the possible samples (data) were collected.

Nitrogen Dioxide

Physical Characteristics and Sources

In its pure state, nitrogen dioxide is a reddish-orangeish-brown gas with a characteristic pungent odor. It is corrosive and a strong oxidizing agent. As a pollutant in ambient air, however, it is virtually colorless and odorless, although it may be an irritant to the eyes and throat. Oxides of nitrogen, nitric oxide and nitrogen dioxide, are formed when the nitrogen and oxygen in the air are combined in high-temperature combustion. Nitric oxide released into ambient air combines with oxygen to form nitrogen dioxide. Major nitrogen dioxide sources are coal conversion processes, natural gas processing plants and natural gas compressor stations.

Health Effects

The negative effects of nitrogen dioxide on personal comfort, well being and the environment include respiratory distress, as well as impacts on vegetation, materials, visibility and acid deposition.⁵ Nitrate aerosols, which result from nitric oxide and nitrogen dioxide combining with water vapor in the air, consistently have been linked to visibility problems.

Standards Comparison

Nitrogen dioxide was monitored at seven sites. Five were operated by the department and two by industry.

The state annual standard (53 ppb) was not exceeded during the year. The maximum annual arithmetic mean of those sites collecting at least 80 percent of the possible data for the year was 6.5 ppb at Fargo NW.

The nitrogen dioxide data are summarized in Table 4.

TABLE 4

COMPARISON OF AIR QUALITY DATA WITH
THE NORTH DAKOTA AMBIENT AIR QUALITY STANDARDS *

POLLUTANT : Nitrogen Dioxide (ppb)

LOCATION	YEAR	SAMPLING PERIOD	NUM OBS	M A X I M A 1 - HOUR		ARITH MEAN	% >MDV
				1ST MM/DD/HH	2ND MM/DD/HH		
Beulah - North	2001	JAN-DEC	8674	40 05/22/16	26 05/22/15	3.4	77.8
DGC #12	2001	JAN-DEC	8570	33 08/26/01	30 07/02/21	3.6	86.1
DGC #17	2001	JAN-DEC	8619	84 11/17/07	66 09/26/06	3.4	76.3
Dunn Center	2001	JAN-DEC	8564	23 07/09/08	22 03/28/20	1.9	44.8
Fargo NW	2001	JAN-DEC	8690	57 03/02/22	56 03/02/20	6.5	81.1
Hannover	2001	JAN-DEC	4737 ***	27 12/06/18	26 02/09/21	2.6	63.5
TRNP - NU	2001	AUG-DEC	3625 ***	9 09/06/22	9 09/07/00	1.3	20.1

The maximum 1-hour concentration is 84 ppb at DGC #17 on 11/17/07

* The air quality standards are:
STATE - 53 ppb maximum annual arithmetic mean.
FEDERAL - 53 ppb annual arithmetic mean.

*** Less than 80% of the possible samples (data) were collected.

Ammonia

Physical Characteristics

Ammonia is a corrosive, colorless gas with a strong irritating odor. It is used in making fertilizer, plastics, dyes, textiles, detergents and pesticides. It reacts with acids and oxidizing materials (fluorine, chlorine, etc.). It is corrosive to copper, zinc and many metal surfaces. It reacts with hypochlorite and halogens to form explosive compounds that are pressure and temperature sensitive.^{6,7}

Health Effects

In mild concentrations (<25 ppm), ammonia will cause conjunctivitis and dermatitis. At higher concentrations, in the eyes it will cause swelling, painful burns, lesions, and possible loss of vision. On contact with the skin, it will cause caustic-like burns and inflammation. Toxic level (300 ppm) skin exposure may cause skin lesions resulting in early necrosis and scarring. Inhalation is corrosive and irritating to the upper respiratory system and all mucus-type tissue. Depending on the concentration inhaled, it may cause burning sensations, coughing, wheezing, shortness of breath, headache and nausea, with eventual collapse and death.^{6,7}

Standards Comparison

There is no ambient air quality standard for ammonia. Because ammonia is important to the newer air quality dispersion models, the ammonia analyzer is maintained at the Beulah - North site.

The ammonia data are summarized in Table 5.

TABLE 5

COMPARISON OF AIR QUALITY DATA WITH
THE NORTH DAKOTA AMBIENT AIR QUALITY STANDARDS *

POLLUTANT : Ammonia (ppb)

LOCATION	YEAR	SAMPLING PERIOD	NUM OBS	M A X I M A					
				1ST	2ND	1 - 3RD	4TH	5TH	6TH
				MM/DD/HH	MM/DD/HH	MM/DD/HH	MM/DD/HH	MM/DD/HH	MM/DD/HH
Beulah - North	2001	JAN-DEC	4737 ***	110.0	91.3	88.3	87.4	76.9	75.1
				05/26/04	11/12/01	05/20/23	07/03/22	02/03/22	12/18/07

*** Less than 80% of the possible samples (data) were collected.

Ozone

Physical Characteristics and Sources

Ozone is a highly reactive form of oxygen. At very high concentrations, it is a blue, unstable gas with a characteristic pungent odor. It often can be detected around an arcing electric motor, lightning storms or other electrical discharges. However, at ambient concentrations, ozone is colorless and odorless.

At ground level where it can be breathed, ozone is a pollutant⁸. However, ground-level ozone should not be confused with the stratospheric ozone located between 12 and 30 miles above the earth's surface. The stratospheric ozone layer shields the earth from intense cancer-causing ultraviolet radiation.⁹ Concentrations of ozone in this layer are approximately 10,000 to 12,000 ppb or one hundred times the SAAQS for ozone. Occasionally, meteorological conditions can result in stratospheric ozone being brought to ground level. This can increase concentrations by 50 to 100 pbb.

Ozone is not emitted directly from a source like other pollutants, but forms as a secondary pollutant. Its precursors are certain hydrocarbons and nitrogen oxides that react chemically in sunlight to form ozone. The sources for these reactive hydrocarbons are automobile exhaust; gasoline and oil storage and transfer; industrial paint solvents; degreasing agents; cleaning fluids; and ink solvents. Nitrogen oxides are created when nitrogen and oxygen in the air combine during high-temperature combustion. Also, vegetation gives off some reactive hydrocarbons; for example, pine trees give off terpene.⁸

Ozone production is a year-round phenomenon. However, the highest ozone levels generally occur during the summer season, when sunlight is stronger and stagnant meteorological conditions can cause reactive pollutants to remain in an area for several days. Ozone produced under these conditions can be transported many miles.

Health Effects

Short-term exposure to ozone in the range of 150 to 250 ppb may impair mechanical functions of the lungs and may induce respiratory difficulties and related symptoms in sensitive individuals (those who have asthma, emphysema or reduced lung function). Symptoms and effects of ozone exposure are more readily induced in people who are exercising.

Ozone is the major component of photochemical “smog,” although the haziness and odors of the smog are caused by other components. The deterioration and degradation of material, especially the splitting and cracking of rubber tires and windshield wiper blades, is associated with ozone. Many plants, such as soybeans and alfalfa, are sensitive to ozone and can be damaged by extended exposure to low levels of ozone.

Standards Comparison

Ozone was monitored at six state-run sites. This data is used in computer dispersion models as part of both the primary and secondary chemical transformation equations.

The 1-hour state standard (120 ppb) was not exceeded during the year. The maximum 1-hour concentration was 70 ppb at Fargo NW.

As part of preparing for a new 8-hour standard (80 ppb), 8-hour averages have been included in the data summary. The 8-hour standard uses the fourth highest daily maximum for comparison to the standard. The highest fourth-highest 8-hour concentration was 63 ppb at Fargo NW.

The ozone data are summarized in Table 6.

TABLE 6

COMPARISON OF AIR QUALITY DATA WITH
THE NORTH DAKOTA AMBIENT AIR QUALITY STANDARDS *

POLLUTANT : Ozone (ppb)

LOCATION	YEAR	SAMPLING PERIOD	NUM OBS	1 - HOUR		M A X I M A 8 - HOUR		1ST 2ND		1ST 2ND 3RD		4TH	1HR #>120	8HR #>80
				MM/DD/HH	MM/DD/HH	MM/DD/HH	MM/DD/HH	MM/DD/HH	MM/DD/HH	MM/DD/HH	MM/DD/HH			
Beulah - North	2001	JAN-DEC	8692	68 04/28/12	68 04/28/13	63 04/28/09	61 04/28/08	61 04/28/10	61 04/28/07					
Dunn Center	2001	JAN-DEC	8703	67 04/28/12	66 05/27/14	63 09/28/11	61 09/28/10	61 09/28/12	61 09/28/09					
Fargo NW	2001	JAN-DEC	8707	70 06/27/17	69 05/19/17	63 06/27/10	63 06/27/11	63 06/28/09	63 05/19/10					
Hannover	2001	JAN-DEC	5475 ***	69 04/28/13	68 04/28/12	65 04/28/08	61 04/28/09	61 04/28/07	61 04/28/10					
TRNP - NU	2001	AUG-DEC	3144 ***	67 09/28/16	66 09/28/14	65 09/28/12	58 09/28/13	58 09/28/11	58 09/28/14					
TRNP - SU (Painted Canyon)	2001	JAN-DEC	8705	64 05/13/14	63 04/28/13	60 09/28/11	59 09/28/12	59 09/28/10	59 09/28/13					

The maximum 1-hour concentration is 70 ppb at Fargo NW on 06/27/17
The 4th highest 8-hour concentration is 63 ppb at Fargo NW on 05/19/10

* The air quality standards for ozone are:
STATE - 120 ppb not to be exceeded more than once per year.

FEDERAL - Fourth highest daily maximum 8-hour averages for a 3-year period not to exceed 80 ppb.

*** Less than 80% of the possible samples (data) were collected

Particulate Matter ($PM_{2.5}$ & PM_{10})

Physical Characteristics and Sources

Particulate matter is the term given to the tiny particles of solid or semi-solid material found in the atmosphere. Particulates ranging in size from less than 0.1 micrometer to 50 micrometers are called Total Suspended Particulate (*TSP*). Particles larger than 50 micrometers tend to settle out of the air quickly and are not considered to have a health impact. Particulate matter 10 micrometers in diameter and smaller is considered inhalable. This particulate matter is called PM_{10} .¹⁰

The majority of anthropogenic (man-made) particulate are in the 0.1 to 10 micrometer diameter range. Particles larger than 10 micrometers usually are due to “fugitive dust” (windblown sand and dirt from roadways, fields and construction sites) and contain large amounts of silica (sand- like) materials. PM_{10} particulate, on the other hand, generally is created during a burning process and includes fly ash (from power plants), carbon black (from automobiles and diesel engines) and soot (from fireplaces and wood-burning stoves). PM_{10} particulates from these sources contain a large percentage of elemental and organic carbon, which play a role in both visual haze and health issues.¹⁰

In addition, particles less than 2.5 micrometers ($PM_{2.5}$) are major contributors to visibility degradation because of their ability to “scatter” light.

Health Effects

The health risk from an inhaled dose of particulate matter depends on the size and concentration of the particulate. Size determines how deeply the inhaled particulate will penetrate into the respiratory tract, where it can persist and cause respiratory damage. Particles less than 10 micrometers in diameter are easily inhaled deeply into the lungs.

Fine particulate ($PM_{2.5}$) pollution affects the health of certain subgroups. Such groups can be identified as potentially “at risk” of adverse health effects from air borne pollutants. There is very strong evidence that asthmatics are much more sensitive (i.e., respond with symptoms at relatively low concentrations) to the effects of particulates than is the general healthy population.⁹

The effects of particulate exposure may be the most widespread of all pollutants. Because of the potential for extremely long-range transport of fine particles and because of the chemical reactions that occur, no place on earth has been spared from the particulate generated by

urban and rural sources. The effects of particulate range from visibility degradation to climate changes to vegetation damage. General soiling, commonly thought to be just a nuisance, can have long-term effects on paint and other materials.¹¹ Acid deposition can be detected in the most remote areas of the world.

Inhalable PM_{2.5} Particulates

Inhalable PM_{2.5} particulates were monitored at seven sites operated by the department. Single-day samplers were installed at Beulah, Dickinson, Sharon, and TRNP - SU to collect a sample once every six days. Sequential samplers were installed at Bismarck, Fargo and Grand Forks to collect a sample once every three days.

Standards Comparison

The 24-hour federal standard ($65 \mu\text{g}/\text{m}^3$) was not exceeded during the year. The maximum 24-hour average concentration was $37.7 \mu\text{g}/\text{m}^3$ at Sharon.

The federal annual standard ($15 \mu\text{g}/\text{m}^3$) was not exceeded for the year. The maximum annual average was $8.1 \mu\text{g}/\text{m}^3$ at Grand Forks - North.

The inhalable PM_{2.5} data are summarized in Table 7.

TABLE 7

COMPARISON OF AIR QUALITY DATA WITH
THE NORTH DAKOTA AMBIENT AIR QUALITY STANDARDS *

POLLUTANT : Inhalable PM_{2.5} Particulates (µg/m³)

LOCATION	YEAR	SAMPLING PERIOD	NUM OBS	MIN	M 1ST MM/DD	A X 2ND MM/DD	I M A 3RD MM/DD	ARITH MEAN	#>150	AM>50	% >MDV
Beulah - North	2001	JAN-DEC	61	1.3	16.4 01/02	16.1 01/12	12.5 01/03	5.8			96.7
Bismarck Residential	2001	JAN-DEC	119	1.8	21.6 01/11	19.7 01/02	17.1 01/04	6.7			98.3
Dickinson Residential	2001	JAN-JUN ***	31	1.8	16.7 01/01	13.0 01/02	12.6 01/02	5.9			93.5
Fargo NW	2001	JAN-DEC	114	0.6	36.0 01/03	25.6 01/04	23.9 01/10	8.2			98.2
Grand Forks - North	2001	JAN-DEC	120	0.5	35.1 01/03	25.9 01/02	22.5 01/10	8.3			96.7
Sharon	2001	JAN-DEC	53	1.6	18.0 01/12	14.1 01/02	12.5 01/04	6.2			98.1
TRNP - SU (Painted Canyon)	2001	JAN-DEC	56	1.5	13.3 01/01	10.9 01/08	10.1 01/02	4.6			91.1

The maximum 24-hour concentration is 36.0 µg/m³ at Fargo NW on 01/03

* The ambient air quality standards are:

FEDERAL Standards -

- 1) 24-hour: 3-year average of 98th percentiles not to exceed 65 µg/m³.
- 2) Annual: 3-year average not to exceed 15µg/m³.

*** Less than 80% of the possible samples (data) were collected.

Inhalable Continuous PM_{2.5} Particulates

Inhalable particulates are monitored continuously at Beulah - North and Fargo NW. Since the data collected is not collected by an EPA reference or equivalent method, the data can not be used for standard comparison. The EPA is expected to designate the analyzers used as an equivalent method some time in 2002.

The maximum 1-hour average concentration was 151.4 µg/m³ at Fargo NW. The maximum 24-hour average concentration is 33.4 µg/m³ at Beulah North. The maximum annual average is 5.6 µg/m³ at Beulah - North.

The inhalable continuous PM_{2.5} data are summarized in Table 8.

Table 8

COMPARISON OF AIR QUALITY DATA WITH
THE NORTH DAKOTA AMBIENT AIR QUALITY STANDARDS *

POLLUTANT : Inhalable Continuous PM_{2.5} (µg/m³)

LOCATION	YEAR	SAMPLING PERIOD	NUM OBS	1 - HOUR		24 - HOUR		3RD	4TH	MEAN	1HR #>150	24HR #>65
				1ST MM/DD/HH	2ND MM/DD/HH	1ST MM/DD	2ND MM/DD					
Beulah - North	2001	JAN-DEC	8718	151.4 03/28/13	150.0 03/28/14	29.7 03/28	17.2 03/24	16.5 02/09	15.4 01/15	6.8	1	
Fargo NW	2001	JAN-DEC	8185	105.5 05/29/21	83.1 08/28/07	25.1 06/28	18.3 07/16	16.4 11/27	16.0 11/06	4.8		

The maximum 1-hour concentration is 151.4 µg/m³ at Beulah - North on 03/28/13
The highest 24-hour concentration is 29.7 µg/m³ at Beulah - North on 03/28

* The ambient air quality standards are:

FEDERAL Standards -

- 1) 24-hour: 3-year average of 98th percentiles not to exceed 65 µg/m³.
- 2) Annual: 3-year average not to exceed 15 µg/m³.

Inhalable PM₁₀ Particulates

Inhalable PM₁₀ particulate concentrations were monitored at three sites.

Standards Comparison

The 24-hour state standard (150 µg/m³) was not exceeded during the year. The maximum 24-hour concentration was 42.4 µg/m³ at Fargo NW.

The annual state standard (50 µg/m³) was not exceeded. The maximum annual mean for the year was 16.8 µg/m³ at Fargo NW.

The inhalable particulate (PM₁₀) data are summarized in Table 9

TABLE 9

COMPARISON OF AIR QUALITY DATA WITH
THE NORTH DAKOTA AMBIENT AIR QUALITY STANDARDS *

POLLUTANT : Inhalable PM₁₀ Particulates (µg/m³)

LOCATION	YEAR	SAMPLING PERIOD	NUM OBS	MIN	M A X I M A			ARITH MEAN	#>150	AM>50	% >MDV
					1ST MM/DD	2ND MM/DD	3RD MM/DD				
Bismarck Residential	2001	JAN-DEC	55	5.6	84.4 05/19	44.1 05/07	41.0 09/28	20.2			100.0
Dragswolf	2001	JAN-DEC	61	0.4	79.0 05/07	37.4 10/04	35.9 10/10	12.7			85.2
White Shield	2001	JAN-DEC	58	0.6	37.2 09/26	34.1 10/04	30.3 10/10	12.2			89.7

The maximum 24-hour concentration is 84.4 µg/m³ at Bismarck Residential on 05/19

* The STATE air quality standards are:

- 1) 150 µg/m³ maximum averaged over a 24-hour period with no more than one expected exceedance per year.
- 2) 50 µg/m³ expected annual arithmetic mean.

SUMMARY AND CONCLUSIONS

The state of North Dakota has relatively clean air. North Dakota is one of only 14 states to comply with all federal ambient air quality standards. The air quality in North Dakota also meets all state ambient air quality standards. Site and pollutant combinations that do not meet the 80 percent data recovery for the full year are reported as a partial year. A summary for each pollutant is provided below.

Sulfur Dioxide

The federal standards were not exceeded at any monitoring site. The state 1-hour standard was exceeded three times. The maximum concentrations and the maximum concentrations expressed as a percentage of the applicable standard are as follows: 1-hour – 164 ppb (60.1%); 3-hour – 116 ppb (23.2%); 24-hour – 38 ppb (38.4%); annual – 5.5 ppb (23.9%).

Sulfur Dioxide 5-Minute Averages

No standard is currently in effect. The maximum 5-minute average was 229 ppb.

Nitrogen Dioxide

Neither state nor federal standard was exceeded at any of the monitoring sites. The maximum concentration and the maximum concentration expressed as a percentage of the applicable standard is as follows: annual – 6.5 ppb (12.3%).

Ammonia

No standard is currently in effect. The maximum 1-hour average is 110.0 ppb.

Ozone

Neither state nor federal standards were exceeded during the year. The maximum 1-hour concentration and the maximum 1-hour concentration expressed as a percentage of the applicable standard is 70 ppb (58.3%). The fourth highest 8-hour average concentration was 63 ppb (78.8%).

Inhalable PM_{2.5} Particulates

The federal standards were not exceeded during the year. The maximum concentrations and the maximum concentrations expressed as a percentage of the applicable standard are as follows: 24-hour – 36.0 $\mu\text{g}/\text{m}^3$ (55.4%); annual – 8.3 $\mu\text{g}/\text{m}^3$ (55.3%).

Inhalable Continuous PM_{2.5} Particulates

No standard is currently in effect for this analytical method. The maximum 1-hour average was 151.4 $\mu\text{g}/\text{m}^3$. The maximum 24-hour average was 29.7 $\mu\text{g}/\text{m}^3$. The maximum annual average was 6.8 $\mu\text{g}/\text{m}^3$.

Inhalable PM₁₀ Particulates

The state standards were not exceeded during the year. The maximum concentrations and the maximum concentrations expressed as a percentage of the applicable standard are as follows: 24-hour – 84.4 $\mu\text{g}/\text{m}^3$ (56.3%); annual – 20.2 $\mu\text{g}/\text{m}^3$ (40.4%).

REFERENCES

REFERENCES

- 1 Environmental Protection Agency, May 1977. Quality Assurance Handbook for Air Pollution Measurement Systems Volume II, Ambient Air Specific Methods (as amended), EPA-600/4-77-027a, Office of Air Quality Planning and Standards, Research Triangle Park, NC.
- 2 Environmental Protection Agency, May 10, 1979. Title 40, Code of Federal Regulations, Part 58 (as amended), United States Government Printing Office, Superintendent of Documents, Washington, DC.
- 3 Environmental Protection Agency, August 7, 1980. Prevention of Significant Deterioration, Title 40, Code of Federal Regulations, Part 52 (as amended), United States Government Printing Office, Washington, DC.
- 4 Environmental Protection Agency, National Air Quality and Emissions Trends Report, 1995, October 1996.
- 5 Environmental Protection Agency Strategies and Air Standards Division, Preliminary Assessment of Health and Welfare Effects Associated with Nitrogen Oxides for Standards-Setting Purposes, U.S. Government Printing Office, Washington; October 1981, pp I-iii.
- 6 New Jersey Department of Health and Senior Services, Hazardous Substance Sheet, Ammonia, June 1998.
- 7 BOC Gases Material Safety Data Sheet, Ammonia (MSDS: G-11), June 1, 1999.
- 8 National Primary and Secondary Ambient Air Quality Standard for Ozone, Title 40 Code of Federal Regulations, Part 50.9 (as amended), United States Government Printing Office, Washington, DC.
- 9 Miller, R. and M. J. Utell, Elements of Meteorology, C. E. Merrill Co., Columbus, Ohio, 1975.
- 10 The Perils of Particulates. American Lung Association, New York, March 1994.
- 11 Sulfur Dioxide, Minimum Lethal Exposure & Maximum Tolerated Exposure, in TOMES Medical Management file [database online]. Colorado Department of Public Health and Environment, 1995 [cited 12 September 1995]. Available from Micromedex Inc. Englewood, Co.

APPENDICES

APPENDIX 1

North Dakota and Federal Ambient Air Quality Standards

STANDARDS

In general, air pollutants are divided into two classes: primary pollutants such as sulfur dioxide, carbon monoxide, nitrogen dioxide, hydrogen sulfide, particulate matter (<2.5 microns) and particulate matter (<10 microns) and secondary pollutants which are formed as the result of a chemical reaction. Sources of primary pollutants include power plants, natural gas processing plants, oil wells, oil refineries, asphalt plants, factories, wind-blown dirt, automobiles, fireplaces and incinerators. Secondary pollutants result from a primary pollutant undergoing a chemical reaction; for example, ozone is formed as a result of a photochemical reaction between hydrocarbons and oxides of nitrogen.

The North Dakota Ambient Air Quality Standards are established to protect public health and welfare. Effective Aug. 1, 1997, coal conversion and oil refineries were exempted from the state sulfur dioxide standards.

Table A1-1 presents the current North Dakota Ambient Air Quality Standards. Table A1-2 presents the federal Ambient Air Quality Standards. State standards must be as stringent as (but may be more stringent than) federal standards.

TABLE A1-1
North Dakota
Ambient Air Quality Standards

Air Contaminants	Standards (Maximum Permissible Concentrations)	
Inhalable Particulate (PM ₁₀)	50	micrograms per cubic meter of air, expected annual arithmetic mean micrograms per cubic meter of air maximum
	150	24-hour average concentration with no more than one expected exceedance per year
Sulfur Dioxide*	0.023	parts per million (60 micrograms per cubic meter of air), maximum annual arithmetic mean concentration
	0.099	parts per million (260 micrograms per cubic meter of air), maximum 24-hour average concentration
	0.273	parts per million (715 micrograms per cubic meter of air), maximum 1-hour average concentration
Hydrogen Sulfide	10.0	parts per million (14 milligrams per cubic meter of air), maximum instantaneous (ceiling) concentration not to be exceeded
	0.20	parts per million (280 micrograms per cubic meter of air), maximum 1-hour average concentration not to be exceeded more than once per month
	0.10	parts per million (140 micrograms per cubic meter of air), maximum 24-hour average concentration not to be exceeded more than once per year
	0.02	parts per million (28 micrograms per cubic meter of air), maximum arithmetic mean concentration averaged over three consecutive months
Carbon Monoxide	9	parts per million (10 milligrams per cubic meter of air), maximum 8-hour concentration not to be exceeded more than once per year
	35	parts per million (40 milligrams per cubic meter of air), maximum 1-hour concentration not to be exceeded more than once per year
Ozone	0.12	parts per million (235 micrograms per cubic meter of air), maximum 1-hour concentration not to be exceeded more than once per year
Nitrogen Dioxide	0.053	parts per million (100 micrograms per cubic meter of air), maximum annual arithmetic mean
Lead	1.5	micrograms per cubic meter of air, maximum arithmetic mean averaged over a calendar quarter

* After Aug. 1, 1997, coal conversion facilities and oil refineries are subject only to the federal SO₂ standards.

TABLE A1-2
Federal Ambient Air Quality Standards

<u>Pollutant</u>	<u>Description</u>	<u>Primary</u>	<u>Secondary</u>
Inhalable Particulate (<2.5 microns)	3-year average of annual arithmetic mean concentrations.	15 $\mu\text{g}/\text{m}^3$	15 $\mu\text{g}/\text{m}^3$
	3-year average of the 98 th percentile of the 24-hour concentrations.	65 $\mu\text{g}/\text{m}^3$	65 $\mu\text{g}/\text{m}^3$
Inhalable Particulates (<10 microns)	Expected annual arithmetic mean.	50 $\mu\text{g}/\text{m}^3$	50 $\mu\text{g}/\text{m}^3$
	99 th percentile of the 24-hour concentrations averaged over 3 years.	150 $\mu\text{g}/\text{m}^3$	150 $\mu\text{g}/\text{m}^3$
Sulfur Dioxide	Annual arithmetic mean.	0.03 ppm (80 $\mu\text{g}/\text{m}^3$)	-
	Maximum 24-hour concentration not to be exceeded more than once per year.	0.14 ppm (365 $\mu\text{g}/\text{m}^3$)	-
	Maximum 3-hour concentration not to be exceeded more than once per year.	-	0.5 ppm (1300 $\mu\text{g}/\text{m}^3$)
Carbon Monoxide	8-hour concentration not to be exceeded more than once per year.	9 ppm (10 $\mu\text{g}/\text{m}^3$)	-
	1-hour average concentration not to be exceeded more than once per year.	35 ppm (40 $\mu\text{g}/\text{m}^3$)	-
Ozone	3-year average of the annual 4 th highest daily maximum 8-hour concentrations, not to be exceeded.	0.08 ppm	0.08 ppm
Nitrogen Dioxide	Annual arithmetic mean.	0.053 ppm (100 $\mu\text{g}/\text{m}^3$)	0.053 ppm (100 $\mu\text{g}/\text{m}^3$)
Lead	Maximum arithmetic mean averaged over a calendar quarter.	1.5 $\mu\text{g}/\text{m}^3$	1.5 $\mu\text{g}/\text{m}^3$


* The PM_{10} and ozone standards have been challenged in court. The final status for these standards is yet to be decided.

APPENDIX 2

Air Quality Personnel Organizational Chart

The following Division of Environmental Engineering organizational chart includes the Air Pollution Control Program.

NORTH DAKOTA DEPARTMENT OF HEALTH
DIVISION OF AIR QUALITY



A2-1 Environmental Engineering Organizational Chart

APPENDIX 3

Wind and Pollution Star Charts

The figures in this appendix are arranged with the site's wind star chart in the upper left-hand position. To remove most of the wind direction bias caused by low wind speeds, wind speeds less than 5 mph were removed from the data. For department-operated sites the pollution star charts are arranged with sulfur dioxide in the upper right-hand position. Next is either hydrogen sulfide or nitrogen dioxide. For industry networks, the wind star chart is presented first followed by the parameters monitored at each site. Except for the Bear Paw - McKenzie Gas Plant network which has wind direction at each site, there is only one MET station for each network.

The pollution star charts present the percentage of time a pollutant is detected when the wind is from a given direction. For example, a wind star chart shows a frequency of 122, and a pollution star chart shows a 66 for the same direction. This means that 66 percent of the time (80 of the possible 122 hours) the wind was greater than 5 mph from that direction and an hourly average for that pollutant had a detectable concentration.

Ozone pollution star charts are not presented because the percentage of time would be essentially 100 percent for each wind sector.






Figure A3-1 Amerada Hess Star Charts








Figure A3-2 Beulah Star Charts







Figure A3-3 Bear Paw Star Charts






Figure A3-4 Dunn Center Star Charts








Figure A3-5 DGC Star Charts





Figure A3-5 DGC Star Charts (cont.)







Figure A3-6 Fargo Star Charts






Figure A3-7 Hannover Star Charts







Figure A3-8 Mandan/Mandan NW Star Charts








Figure A3-9 TRNP - NU Star Charts


TRNP - SU (Painted Canyon) Wind Direction Star Chart during 2001



Percent of Time SO₂ Detected for a Given Wind Sector for TRNP - SU (Painted Canyon) during 2001



White Shield Wind Direction Star Chart during 2001



Percent of Time SO₂ Detected for a Given Wind Sector for White Shield during 2001




Figure A3-10 Short Creek - SPM/TRNP - NU

APPENDIX 4

1992-2001 Trends

The trend graphs for 1992 through 2001 are presented in alphabetical order, grouped by site, unless multiple sites would fit on a single page. Each graph depicts the maximum concentration for each applicable standard (left scale) and percentage of time an hourly concentration is detected (right scale).









Figure A4-1 Amerada Hess/Beulah North Trends









Figure A4-2 Beulah North (cont.)/Bear Paw Trends









Figure A4-3 DGC Trends









Figure A4-4 Dunn Center/Dickinson/Dragswolf Trends














Figure A4-5 Fargo NW Trends



A4-6 Grand Forks - North/Hannover Trends







Figure A4-7 Mandan/Mandan NW Trends






Figure A4-8 TRNP - NU Trends







Figure A4-9 TRNP - SU Trends

