

# Nutrient Characteristics for Streams in North Dakota

**By Joel M. Galloway**  
**Associate Director**

**USGS North Dakota Water Science Center**



# Background

- Study completed in 2012 examined data from various agencies that have conducted water-quality sampling programs and projects for streams in ND for a number of years for various purposes.
- The purpose of the study was to:
  1. Provide descriptive statistics and summaries of water-quality data from sites throughout the State;
  2. Determine trends and loads for selected constituents and sites with sufficient concentration and streamflow data;
  3. Determine an efficient state-wide network sampling design for monitoring future water-quality conditions



# Nutrient Data and Analysis

- Examined all data available from 1970-2008 -  
Data obtained from USGS NWIS database, USEPA STORET database, and electronic files from NDDH
- Data screening yielded 186 sites across the State with 10 or more samples for most constituents
  - Nutrients – 50,880 values after data was processed


# Nutrient Characteristics

- Spatial distribution of concentrations
- Yields
- Trends





# SPATIAL DISTRIBUTION - MEDIAN AMMONIA CONCENTRATIONS



Base from U.S. Geological Survey digital data, 1983  
Universal Transverse Mercator projection Zone 14


## EXPLANATION

Concentration, in milligrams per liter as nitrogen

- ◆ Less than 0.04
- ◆ 0.04 to 0.06
- ◆ 0.06 to 0.09
- ◆ Greater than 0.09

**64 of 139 sites – median <0.04 mg/L**  
**Greatest median conc. – 0.59 mg/L**


# SPATIAL DISTRIBUTION - MEDIAN NITRATE PLUS NITRITE CONCENTRATIONS






# SPATIAL DISTRIBUTION - MEDIAN DISSOLVED PHOSPHORUS CONCENTRATIONS


Period of record




Spring



Summer



Winter



Base from U.S. Geological Survey digital data, 1983  
Universal Transverse Mercator projection Zone 14





## EXPLANATION

Concentration, in milligrams per liter as phosphorus

- ◆ Less than 0.01 to 0.03
- ◆ 0.03 to 0.09
- ◆ 0.09 to 0.16
- ◆ Greater than 0.16

11 of 128 sites – median <0.01 mg/L  
Greatest median conc. – 0.55 mg/L


# SPATIAL DISTRIBUTION - MEDIAN TOTAL PHOSPHORUS CONCENTRATIONS



Base from U.S. Geological Survey digital data, 1983  
Universal Transverse Mercator projection Zone 14


## EXPLANATION

Concentration, in milligrams per liter as phosphorus












1 of 128 sites – median <0.01 mg/L  
Greatest median conc. – 0.51 mg/L





From Mueller and Spahr (2006)

**Figure 2.** Location of the 481 National Water-Quality Assessment sampling sites with adequate data for analysis of nutrient concentrations and loads.



From Mueller and Spahr (2006)

**Figure 14.** Box plots showing the distribution of flow-weighted nutrient concentrations by land-use category.



# Loads and Yields

- Loads and Yields were estimated for nitrogen and phosphorus
- 34 sites were selected with sufficient data to determine reasonable estimates of load
- Loads (lbs/yr) were estimated using LOADEST to determine regression models:

$$\ln(\text{Load}) = \beta_0 + \beta_1 \ln(\text{Daily Streamflow}) + \beta_2 (\text{Time}) + \beta_3 \sin(2\pi \text{Time}) + \beta_4 \cos(2\pi \text{Time})$$


**Seasonality**

- Yields (lbs/yr/mi<sup>2</sup>) were calculated from the estimated loads:

$$\text{Load (lbs/yr)} / \text{Drainage area (mi}^2\text{)}$$

# Nutrient Yields

Nitrate plus nitrite (total and dissolved)




## EXPLANATION

Normalized annual yields, in pounds per year per square mile



Ammonia (total and dissolved)




## EXPLANATION

Normalized annual yields, in pounds per year per square mile



Dissolved phosphorus




## EXPLANATION

Normalized annual yields, in pounds per year per square mile



Total phosphorus



## EXPLANATION

Normalized annual yields, in pounds per year per square mile



## Nitrate plus Nitrite

MRB:

25 to 120 lbs/yr/mi<sup>2</sup>

RRB:

2 to 1,260 lbs/yr/mi<sup>2</sup>

## Ammonia

MRB:

6 to 22 lbs/yr/mi<sup>2</sup>

RRB:

4 to 62 lbs/yr/mi<sup>2</sup>

## Dissolved phos.

MRB:

<1 to 35 lbs/yr/mi<sup>2</sup>

RRB:

5 to 47 lbs/yr/mi<sup>2</sup>

## Total phosphorus

MRB:

<1 to 167 lbs/yr/mi<sup>2</sup>

RRB:



6 to 76 lbs/yr/mi<sup>2</sup>



# Trends



- Time-series model (QWTREND) used to evaluate flow-related variability and trends in historical concentrations. Significant trends were determined using maximum likelihood estimation and generalized likelihood ratio tests
- Selected 10 sites with sufficient data
  - Sites had to have:
    - » At least 15 years of data
    - » At least 4 samples per year
    - » Have both major ion and nutrient data
- Trends were evaluated for nitrate plus nitrite and total phosphorus

# Nitrate plus Nitrite Trends





# Total Phosphorus Trends



# Summary

- Spatial Distribution:
  - Ammonia conc. higher in winter
    - No distinct spatial pattern
  - Nitrate plus nitrite - higher conc. in Red River Basin
    - Higher in winter in the Missouri Basin, lower in winter in Red River basin
  - Diss. phosphorus - higher in Red River Basin
    - No distinct seasonal pattern
  - Total phosphorus - higher in Red River Basin
    - Highest in summer
  - Total organic carbon – higher in Missouri Basin
- Nutrient yields generally higher in Red River Basin
- Nutrient Trends:
  - Nitrate plus Nitrite – no trend at most sites, upward trend in Red River
  - Total phosphorus – 4 sites had downward trend, 1 site had upward trend (Red River), others had no trend

# ANY QUESTIONS?

Data and analysis are from:

## **Evaluation of Water-Quality Characteristics and Sampling Design for Streams in North Dakota, 1970–2008**

By Joel M. Galloway, Aldo V. Vecchia, Kevin C.  
Vining, Brenda K. Densmore, and  
Robert F. Lundgren

<http://pubs.usgs.gov/sir/2012/5216/>

